

The Miller School of
Medicine at the
University of Miami
Miami, FL 33136

Health Education
International
1530 South Shore Rd.
Ocean View, NJ 08230

References

- Berger, V., Perier, S., Pachiaudi, C., Normand, S., Louisot, P., and Martin, A. (1998) Dietary specific sugars for serum protein enzymatic glycosylation in man. *Metabolism*, 47, 1499-1503.
- t'Hart, L.A., van den Berg, A.J., Kuis, L., van Dijk, H., and Labadie, R.P. (1989) An anti-complementary polysaccharide with immunological adjuvant activity from the leaf parenchyma gel of Aloe vera. *Planta Med.*, 55, 509-512.
- Murray, R.K., (2000) Glycoproteins. In Murray RK, Granner DK, Mayes PA, Rodwell VW (eds): *Harpers Biochemistry*.
- Axford, J. (1997) Glycobiology and medicine: an introduction. *J. R. Soc. Med.*, 90, 260-264.
- Sharon, N. and Lis, H. (1993) Carbohydrates in cell recognition. *Sci. Am.*, 268, 82-89.
- Kelly, G.S. (1999) Larch arabinogalactan: clinical relevance of a novel immune-enhancing polysaccharide. *Altern. Med. Rev.*, 4, 96-103.
- Melikian, H.E., Ramamoorthy, S., Tate, C.G., and Blakely, R.D. (1996) Inability to N-glycosylate the human norepinephrine transporter reduces protein stability, surface trafficking, and transport activity but not ligand recognition. *Mol. Pharmacol.*, 50, 266-276.
- Taylor, M. (2006) *Introduction to Glycobiology*.
- Bastian, W., Zhu, J., Way, B., Lockwood, D., and Livingston, J. (1993) Glycosylation of Asn397 or Asn418 is required for normal insulin receptor biosynthesis and processing. *Diabetes*, 42, 966-974.
- Kanoh, T., Saito, K., Matsunaga, K., Oguchi, Y., Taniguchi, N., Endoh, H., Yoshimura, M., Fujii, T., and Yoshikumi, C. (1994) Enhancement of the antitumor effect by the concurrent use of a monoclonal antibody and the protein-bound polysaccharide PSK in mice bearing a human cancer cell line. *In Vivo*, 8, 241-245.
- Lassalle, B. and Testart, J. (1996) Lectins binding on human sperm surface increase membrane permeability and stimulate acrosomal exocytosis. *Mol. Hum. Reprod.*, 2, 651-658.
- Deal, C.L. and Moskowitz, R.W. (1999) Nutraceuticals as therapeutic agents in osteoarthritis. The role of glucosamine, chondroitin sulfate, and collagen hydrolysate. *Rheum. Dis. Clin. North Am.*, 25, 379-395.
- Choi, B.H., Yee, S., and Robles, M. (1996) The effects of glutathione glycoside in methyl mercury poisoning. *Toxicol. Appl. Pharmacol.*, 141, 357-364.
- Salyers, A.A. (1985) Breakdown of polysaccharides by human intestinal bacteria. *J. Environ. Pathol. Toxicol. Oncol.*, 5, 211-231.
- Pulverer, G., Beuth, J., and Ko, H.L. (1994) Importance of lectins for the prevention of bacterial infections and cancer metastases. *Zentralbl. Bakteriol.*, 281, 324-333.
- David, G. (1992) Structural and functional diversity of the heparan sulfate proteoglycans. *Adv. Exp. Med. Biol.*, 313, 69-78.
- Warczynski, P., Gil, J., Szmigielski, S., Beuth, J., and Pulverer, G. (1997) Prevention of hepatic metastases by liver lectin blocking with D-galactose in colon cancer patients. A prospectively randomized clinical trial. *Anticancer Res.*, 17, 1223-1226.
- Niehues, R., Hasilik, M., Alton, G., Korner, C., Schiebe-Sukumar, M., Koch, H.G., Zimmer, K.P., Wu, R., Harms, E., Reiter, K., von Figura, K., Freeze, H.H., Harms, H.K., and Marquardt, T. (1998) Carbohydrate-deficient glycoprotein syndrome type Ib. Phosphomannose isomerase deficiency and mannose therapy. *J. Clin. Invest.*, 101, 1414-1420.
- Tomana, M., Schrohenloher, R.E., Koopman, W.J., Alarcon, G.S., and Paul, W.A. (1988) Abnormal glycosylation of serum IgG from patients with chronic inflammatory diseases. *Arthritis Rheum.*, 31, 333-338.
- Tomana, M., Schrohenloher, R.E., Reveille, J.D., Arnett, F.C., and Koopman, W.J. (1992) Abnormal galactosylation of serum IgG in patients with systemic lupus erythematosus and members of families with high frequency of autoimmune diseases. *Rheumatol. Int.*, 12, 191-194.
- Wiese, T.J., Dunlap, J.A., and Yorek, M.A. (1997) Effect of L-fucose and D-glucose concentration on L-fucoprotein metabolism in human Hep G2 cells and changes in fucosyltransferase and alpha-L-fucosidase activity in liver of diabetic rats. *Biochim. Biophys. Acta*, 1335, 61-72.
- Tertov, V.V., Orekhov, A.N., Sobenin, I.A., Morrisett, J.D., Gotto, A.M., Jr, and Guevara, J.G., Jr (1993) Carbohydrate composition of protein and lipid components in sialic acid-rich and -poor low density lipoproteins from subjects with and without coronary artery disease. *J. Lipid Res.*, 34, 365-375.
- Kai, H., Murata, Y., Ishii, T., Nishijima, S., Murahara, K., Ogasawara, S., Sugiyama, N., Takahama, K., and Miyata, T. (1990) Anti-allergic effect of N-acetyl neuraminic acid in guinea-pigs. *J. Pharm. Pharmacol.*, 42, 773-777.
- Murch, S.H., MacDonald, T.T., Walker-Smith, J.A., Levin, M., Lionetti, P., and Klein, N.J. (1993) Disruption of sulphated glycosaminoglycans in intestinal inflammation. *Lancet*, 341, 711-714.
- Turner, M.W. (1998) Mannose-binding lectin (MBL) in health and disease. *Immunobiology*, 199, 327-339.

Division of Complementary Medicine
MILLER | SCHOOL OF MEDICINE
UNIVERSITY OF MIAMI

Postage

This Review is
published jointly
by the Division of
Complementary
Medicine at the
University of
Miami's Miller
School of Medi-
cine and Health
Education
International

Advisory Board:

Mahendra Kumar, Ph.D.
Diego A. Deleon, M.D.
Jay K. Ober, ARNP
Francis Quinones, Pharm D.

April, 2007

Inside This Issue:

- History
- Synopsis
- Causes
- Disease Implications

VOLUME 1
ISSUE 1

GLYCOBIOLOGY review

The Language of Life

an overview and history of the science of glycobiology

Antiquated Thinking

It is widely taught in educational institutions around the world that there are four major classes of biomolecules: proteins, fats, nucleic acids, and carbohydrates. These molecules have specific and limited uses in the body, and if the diet contains the correct proportions, coupled with minor amounts of vitamins and minerals, proper nutrition is achieved.

Research from the last few decades, however, has exposed limitations in this traditional view, particularly in the role of carbohydrates (sugars). It is almost universally accepted that carbohydrates play a role solely in the production of energy. It is becoming increasingly apparent that this theory is inaccurate, as sugars play a far more complex role than originally thought.¹ This expanded view has led to a new genre of science called Glycobiology, an area of research with exciting possibility.

History of Glycobiology

When the body is breached by microscopic invaders, how does the immune system know foreigners from healthy cells? What leads the digestive tract to absorb nutrients and repel bacteria? What changes when the body suddenly begins attacking itself in autoimmune disease or fails to defend the body in infections and cancer? The answers to these questions are critical to maintaining health.


In the Cold War era, following World War II, interest in the Aloe Vera plant became widespread as it was discovered that the plant had therapeutic value in its ability to heal. Despite its inclusion in a wide variety of products, it took 30 years to discover the reason why this plant had such beneficial proper-

ties. Researchers finally identified a single monosaccharide, Mannose,² as the therapeutic molecule in this plant and the science of Glycobiology gained momentum.

The Discovery

This discovery of health benefits from a little known monosaccharide led to the search for other sugars of potential value. What was eventually discovered was the fact that eight individual sugars are utilized by the body for the synthesis of glycoconjugates, molecules imperative to the health of many body systems.³

The magnitude of this discovery was not immediately recognized due to the persistent belief that saccharides are used only for energy production. This limited view still persists today despite a growing body of contradictory evidence. Multiple studies have determined that each of the saccharides have specific roles in the body beyond mere energy production.⁴ These tasks include, but are not limited to, cellular communication, structural integrity, physical protection, and cellular adhesion. Because of the critical nature of these saccharides, it is alarming to note that many people suffer monosaccharide deficiencies due to a modern diet lacking in these crucial nutrients.



An Overview and Review of the Science of Glycobiology

Cellular Communication

Monosaccharides are utilized by the body for cell to cell communication. These sugars are arranged in various combinations and serve as an intercellular Braille. Healthy cells exhibit specific combinations that are recognized by other body cells, while foreign combinations of saccharides are not recognized and are tagged for elimination.⁵ Correct configurations of monosaccharides are especially vital for immune system cellular communication, as it is the job of these specialty cells to recognize and eliminate foreign invaders.^{6,25}

Communication is also vital for proper nerve transmission since receptors in the synaptic cleft have specific sugar configurations that are only activated by neurotransmitters when constructed properly.⁷

Red blood cells are coated in glycoproteins with specific sugar configurations as well. In fact, exact sugar combinations are so vital that changing a single sugar means the difference between Type-A and Type-B blood.⁸

Other examples that illustrate the importance of these saccharides to intercellular communication are insulin receptors,⁹ antibodies recognizing and binding with pathogens,¹⁰ and the joining of a sperm cell and the ovum in fertilization.¹¹

Structure

Sugar conjugates not only assist the cell in communication but are also integral to the stability of many body components. Connective tissue is highly dependent upon various saccharide building blocks.⁸ Proteoglycans (polysaccharide gels) are a major component of connective tissues and, besides acting as tissue glue, serve to slow the spread of pathogens. Glycosaminoglycans (GAGs) are polysaccharide structures that possess negative charges which allow them to attract sodium and potassium, thereby attracting water. This absorption plays a key role in water and electrolyte balance, as well as serving as a protective cushion against external forces. Chondroitin, the most abundant GAG, is an important building block of cartilage and is

Structure

highly dependent upon dietary saccharides for proper construction.¹²

The breakdown of nutrients and exposure to the environment causes the production of free radicals that can damage healthy tissue and genetic material. These free radicals are combated by antioxidants, the most important being glutathione. This molecule is constructed and utilized more effectively in the presence of certain saccharides.¹³

The delivery of many components depends upon transport proteins, and it is the saccharide portion of these trafficking proteins that both binds the package and determines its destination.⁸

Protection

The location of carbohydrates as the outer molecule on cell surfaces allows them to act as a cellular barrier from the environment. This role is seen in the mucosal lining of the gastrointestinal tract, where saccharide molecules block the adhesion of toxins and foreign matter while allowing nutrients and harmless molecules to bind and absorb.¹⁴

When properly constructed, cell membrane saccharides can competitively block virus and bacteria from binding and penetrating the cell.¹⁵ These saccharides are also utilized in the circulatory system where they bind directly to foreign pathogens, preventing the penetration of these foreigners.

Adhesion

Saccharide conjugates are utilized in cellular adhesion. This has direct implications for platelet aggregation and tissue stability. A certain saccharide conjugate called heparin is an important molecule for halting the clotting cascade.¹⁶ Deficiencies in construction of this molecule can lead to hypercoagulation, which

in turn can result in heart attack or stroke.

As previously discussed, tissue stability is mainly a component of the saccharide glue and intercellular saccharide molecule linkages. Defects in this area are being investigated as a possible source of cancer metastasis, since malignant tumors often lose their adhesive properties and migrate to other parts of the body.^{15,17}

Saccharide Deficiencies

Almost every cell of the body depends upon these saccharides to function properly, and yet the modern diet is woefully deficient in these vital molecules. To counteract this lack of access to dietary saccharides, the body utilizes a backup system to generate any of the saccharides from glucose. This process depends upon multiple enzymatic steps and excess energy to occur.

A growing body of research suggests that a breakdown in this process is the cause of many common pathologies.¹ If a breakdown does occur, and dietary saccharides are not available, deficiencies will occur.

The most common causes of breakdowns in enzymatically constructing monosaccharides are vitamin deficiencies, stress, illness and aging. Also, genetic errors of saccharide metabolism

result in the deficiency of at least one of the monosaccharides. It is important to note that being born with an inability to generate even one monosaccharide is fatal, unless prompt supplementation of that missing nutrient is supplied.¹⁸ This illustrates the importance of these saccharides and the dire consequences of even minor deficiencies.


Disease Implications

Deficiencies or alterations in monosaccharide conjugates are the suspected culprits in a large number of pathologies, including some

“An inability to generate even one monosaccharide is fatal...”

Saccharides Required for Glycoprotein Synthesis:

Glucose (Glu)
Galactose (Gal)
Mannose (Man)
Fucose (Fuc)
Xylose (Xy)
N- Acetylglucosamine (GlcNAc)
N- Acetylgalactosamine (GalNAc)
N- Acetylneuraminic Acid (NANA)



of the top killers in America.

Various inflammation disorders have been studied in relation to saccharides, including Rheumatoid Arthritis and Lupus. The sugars are imperative in their role as communication molecules on the binding end of the antibody IgG, which may signal false and prolonged inflammation of connective tissues.^{19,20}

Certain autoimmune diseases are believed to be linked to saccharides, including Diabetes Mellitus Type-1 and Multiple Sclerosis. The common factor in these conditions is an error in glycosylation, or the construction of glycoconjugates, which can lead to an attack of healthy tissue by immune cells.²¹

Saccharides may also play a key role in the pathology of certain cancers. Even a healthy human body can produce numerous cancerous cells on a daily basis, but these are typically disposed of by the immune system. Only when immune system fails to recognize the mutated cells do cancers begin their relentless growth to mature tumors. Improper saccharide construction has been investigated as a possible cause of the failure of immune cells to recognize cancer.¹⁰

Other pathologic conditions that have been studied in connection with saccharide deficiencies or alterations include cardiovascular disease²², hypersensitivity, allergies²³, ulcerations of the gastrointestinal and genitourinary tract²⁴, infertility¹¹, neurological and psychiatric conditions.⁷ Future issues of this publication will focus on many of these pathologies in detail.

Chad Reihm - University of Miami